



## Egypt

The following information must be viewed as a guide only. It is not intended, nor implied to be a substitute for professional medical advice.

Specific recommendations on vaccinations, antimalarial medications & targeted travel health advice is always provided on an individual basis taking into account:

- the personal health of the traveller including past medical & vaccination history;
- intended activities;
- precise itinerary;
- style of travel;
- type of accommodation;
- time of year;
- altitude; &
- length of stay.

As well, some vaccines eg rabies & tuberculosis are very much influenced by local disease risk. Specific face-to-face advice is particularly important when recommending antimalarial medications & those for presumptive treatment eg for travellers diarrhoea.

We strongly recommend travellers seek an appointment with a doctor trained in travel health prior to departure.

Medical & nursing staff at The Travel Doctor-TMVC are trained in international public health issues with a focus on immunisations & preventive medicine. Many have travelled extensively & a number have worked in less developed areas of the world for extended periods. Travellers should undergo individual risk assessments whether they are short term holiday makers, business people or the long term expatriate worker. The Travel Doctor-TMVC has clinics Australia wide. In Australia the local centre may be contacted on 1300 658 844, or by visiting [www.traveldoctor.com.au](http://www.traveldoctor.com.au). It is recommended that you visit a travel health professional 6-8 weeks prior to departure. However, if that time frame is not possible, remember – “It’s never too late to vaccinate”.

### Healthy Travelling in Egypt

Egypt is becoming a major destination for Australian travellers. The magnificence of the pyramids & other wonders of ancient Egypt together with the natural beauty of the Nile are enduring drawcards.


Pre-travel preparation will help protect your health while you are away. To assist you in recognising & understanding some of the major travel health risks you may face while holidaying in Egypt, The Travel Doctor-TMVC has prepared a summary of some of these issues in the following pages.

**Table 1** provides a brief description of some of the major travel health issues & vaccinations that should be considered for travel to Egypt.

**Table 2** provides a summary of these major travel health issues & preventative measures that should be considered.


We hope you find this information useful in preparing for your trip. Remember it is advisable to consult a travel health specialist prior to departure.

Currency of your basic immunisations such as Tetanus & Diphtheria should be checked & others like Hepatitis A & Typhoid considered according to the criteria mentioned previously.



**Table 1. Major Travel Health Issues & Considerations for Egypt**

Hepatitis A	This is a viral disease of the liver which is transmitted through eating contaminated food or drinking contaminated water. It is the most common vaccine preventable disease that occurs in travellers to less developed areas of the world. It is strongly recommended for travel to Egypt.
Hepatitis B	This is a viral disease of the liver that is transmitted via blood, blood products or bodily fluids. It is vaccine preventable. Hepatitis B immunisation is now part of the childhood immunisation schedule. Many adult travellers have missed this very important immunisation & travel may be a good reason for vaccination. The vaccine is currently provided to all children as part of our childhood immunisation program.
Typhoid	Typhoid Fever is caused by a bacteria found in contaminated food & water. It is endemic in the developing world & vaccination is recommended for travellers to areas where environmental sanitation & personal hygiene may be poor. The adventurous eater venturing 'off the beaten' path should certainly consider vaccination.
Tetanus, Pertussis & Diphtheria	Tetanus is caused by a toxin released by a common dust or soil bacteria, which enters the body through a wound. Diphtheria is a bacterial infection of the throat & occasionally of the skin. It is found world wide & is transmitted from person-to-person by coughing & sneezing. Pertussis (Whooping cough) is a highly infectious respiratory infection responsible for 300,000 deaths annually, mainly in children. Diphtheria & pertussis vaccines can be added to the tetanus vaccine. Because many adults no longer have immunity from childhood immunisation it is advised that travellers to less developed countries have a tetanus, diphtheria & pertussis booster.
Measles, Mumps & Rubella	Childhood immunisation coverage in many developing countries is not good. As such, travellers under the age of 40 years should have their measles, mumps & rubella immunisation complete. Those over the age of 40 years are most likely to have long term immunity from previous exposure as a child.
Chickenpox	This very common infectious disease can now be prevented through immunisation. Many people miss the disease in childhood only to have a significant illness as an adult. Travel puts one at higher exposure & if one cannot elicit a history of having had the illness a test can show whether at risk.
Influenza	Individuals intending to travel out of an Australian winter might consider the current flu vaccine at the beginning of the season. Exposure to illness in airports & commuter transport is common & exposure may ruin a much needed break. In fact, influenza is likely to be the commonest vaccine preventable disease faced by travellers.
Poliomyelitis	All travellers to Egypt should consider vaccination against polio. Poliomyelitis is a viral infection that can lead to paralysis & sometimes death. Transmission is by faecal contamination of food, usually by unhygienic food handlers or flies, or directly from infected nasal secretions. Although most Australian's & New Zealanders will have been immunised in childhood, it is important to note that efficacy wanes after 10 years & a booster dose is recommended if travelling to a country where the disease could still found, such as Egypt. Polio was still considered endemic in Egypt until recently, although there has been no reported cases since May 2004.
Malaria	Malaria is transmitted by a night biting mosquito. The risk of exposure is low in Egypt, especially in the areas frequented by travellers. While the risk is very low it is not NIL & discussing your particular trip with a travel medicine specialist is important. Medications to reduce the risk of disease are not generally recommended but any illness which is flu-like in nature after returning should still be checked for malaria.
Rabies	Rabies is a deadly viral infection of the brain transmitted to humans. The disease itself is rare in travellers, but the risk increases with extended travel & the likelihood of animal contact. The best way to avoid rabies is to avoid all contact with animals. Dogs are the main carriers; however monkeys, bats, cats & other animals may also transmit the disease. Pre-exposure vaccination is recommended for extended travel & those who work with, or are likely to come in contact with animals.
Cholera	Cholera is a severe, infectious diarrhoeal disease caused by a bacteria. It is common in developing countries & is associated with conditions of poverty & poor sanitation. Cholera causes severe & rapid dehydration. Travellers who follow the rules of eating & drinking safely will minimise their risk. There is also a new oral vaccine available for cholera which may be recommended under certain circumstances.
Traveller's Diarrhoea	Up to 40% of tourists may develop 3 or more loose bowel motions a day within the first week of travel. A variety of germs can be responsible for this infection & a Traveller's Medical Kit containing appropriate therapy can rapidly improve the symptoms. It is also important to follow the rules of healthy eating & drinking to minimise risks.





**Table 2. Summary of travel health issues for Egypt & preventative options available**


	Vaccine Available	Healthy Eating & Drinking	Insect Avoidance	Animal Avoidance
Hepatitis A	✓	✓		
Hepatitis B	✓			
Typhoid	✓	✓		
Tetanus, Pertussis & Diphtheria	✓			
Measles, Mumps & Rubella	✓			
Chickenpox	✓			
Influenza	✓			
Poliomyelitis	✓			
Malaria			✓	
Rabies	✓			✓
Cholera	✓	✓		
Traveller's Diarrhoea		✓		

✓ indicates preventative considerations

**Yellow Fever** vaccination may be required for all travellers arriving from or transiting through Yellow Fever infected areas, such as Central Africa or the Americas.



Remember to check the DFAT 'Smartraveller' website [www.smartraveller.gov.au](http://www.smartraveller.gov.au) prior to departure



...for a safe and healthy journey

Travel Health Fact Sheet **Egypt**

Copyright © The Travel Doctor-TMVC Pty Ltd 2007

Clinics Australia Wide. For your nearest clinic call **1300 658 844** or visit [www.traveldoctor.com.au](http://www.traveldoctor.com.au)



The Travel Doctor-TMVC is proud to be an official supplier to the ARU & the Wallabies